

Kemit Medical Group



23077 Greenfield RD. STE 231

Southfield, MI 48075

SuPrep Bowel Prep for PM Colonoscopy

Carton Contains:

2- 6 ounce bottles of liquid bowel prep

1- 16 ounce mixing container

The DAY BEFORE your Colonoscopy: Clear liquids only

Absolutely No Solid Foods

Examples of clear liquids: Water, clear fruit juices such as apple or white grape, chicken or beef bouillon, jello (no RED or PURPLE), clear Gatorade, popsicles (no RED or PURPLE), clear soft drinks, coffee without cream or sugar.

NO MILK OR MILK PRODUCTS, NO ORANGE JUICE. NO RED OR PURPLE JELLO OR JUICES.

5PM: Pour ONE 6-ounce bottle of SUPREP liquid into the mixing container. Add water to the mixing container up to the red fill line. Drink all the liquid in the container at once. Drink two glasses of water, using the mixing container once again filling the water up to the red fill line over the hour after drinking the solution.

NOTHING BY MOUTH AFTER MIDNIGHT.

The DAY OF your Colonoscopy:

6AM: Pour ONE 6-ounce bottle of SUPREP liquid into the mixing container. Add water to the mixing container up to the red fill line. Drink all the liquid in the container at once. Drink two glasses of water, using the mixing container once again filling the water up to the red fill line over the hour after drinking the solution.

REMEMBER: The preparation is very important. An adequate clean out allows for the best evaluation of your entire colon. During the prep, using baby wipes may ease some of your discomfort. You should NOT plan on working or driving on the day of the procedure due to sedation given at the procedure.

No aspirin, aspirin by-products or plavix for 1 week prior to your colonoscopy. No Coumadin for 5 days or check with your physician who orders the coumadin. Please contact the physician that prescribed the aspirin, plavix, and coumadin to see if this is acceptable.

Kemit Medical Group



23077 Grennfield RD. STE 231
Southfield, MI 48075

Date: _____

Ms/Mr. _____

You are scheduled for a _____ on

Date: _____ Time: _____ at _____ Hospital.

For any date change or cancellation please notify the office at least 24 hours in advance. If we are not notified you will be subject to a \$50 cancellation fee. If a referral is required, you must obtain a referral from your primary care physician office. You must have your referral faxed to our office at least 3 days before your procedure date. If a referral is not received at that time, the procedure will be cancelled. The referral used for your office visit cannot be used for an outpatient or radiological procedure.

Thank You,

Dr. A. D. Williams and Staff

Kemit Medical Group



23077 Greenfield RD. STE 231

Southfield, MI 48075

SuPrep Bowel Prep for AM Colonoscopy

Carton Contains:

2- 6 ounce bottles of liquid bowel prep

1- 16 ounce mixing container

The DAY BEFORE your Colonoscopy: Clear liquids only

Absolutely No Solid Foods

Examples of clear liquids: Water, clear fruit juices such as apple or white grape, chicken or beef bouillon, jello (no RED or PURPLE), clear Gatorade, popsicles (no RED or PURPLE), clear soft drinks, coffee without cream or sugar.

NO MILK OR MILK PRODUCTS, NO ORANGE JUICE. NO RED OR PURPLE JELLO OR JUICES.

3PM: Pour ONE 6-ounce bottle of SUPREP liquid into the mixing container. Add water to the mixing container up to the red fill line. Drink all the liquid in the container at once. Drink two glasses of water, using the mixing container once again filling the water up to the red fill line over the hour after drinking the solution.

7PM: Pour ONE 6-ounce bottle of SUPREP liquid into the mixing container. Add water to the mixing container up to the red fill line. Drink all the liquid in the container at once. Drink two glasses of water, using the mixing container once again filling the water up to the red fill line over the hour after drinking the solution.

NOTHING BY MOUTH AFTER MIDNIGHT AND THE DAY OF THE PROCEDURE.

REMEMBER: The preparation is very important. An adequate clean out allows for the best evaluation of your entire colon. During the prep, using baby wipes may ease some of your discomfort. You should NOT plan on working or driving on the day of the procedure due to sedation given at the procedure.

No aspirin, aspirin by-products or plavix for 1 week prior to your colonoscopy. No Coumadin for 5 days or check with your physician who orders the coumadin. Please contact the physician that prescribed the aspirin, plavix, and coumadin to see if this is acceptable.

Kemit Medical Group



23077 Grennfield RD. STE 231
Southfield, MI 48075